

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : A61K 35/78, C07H 15/26, 3/04		A1	(11) International Publication Number: WO 99/34812 (43) International Publication Date: 15 July 1999 (15.07.99)
(21) International Application Number: PCT/US99/00334 (22) International Filing Date: 7 January 1999 (07.01.99) (30) Priority Data: 09/005,934 12 January 1998 (12.01.98) US (71)(72) Applicant and Inventor: XU, Kai, Jian [CN/US]; Unit 604, 901 S. Plymouth Court, Chicago, IL 60605 (US). (74) Agents: PILLAI, Xavier, Leydig, Voit & Mayer, LTD., Suite 300, 700 Thirteenth Street, N. W., Washington, DC 20005-3960 (US) et al.			(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.
(54) Title: PHARMACEUTICAL COMPOSITIONS AND METHOD OF USING SAME			
(57) Abstract <p>The present invention provides a pharmaceutical composition comprising an extract or combination of extracts having an antiviral, antibacterial, or immunomodulating property, wherein the extract or combination of extracts is obtained from a combination of plants wherein at least one plant is selected from each of the genres <i>Labiatae</i>, <i>Caprifoliaceae</i>, and <i>Oleaceae</i>. A preferred extract is obtained from <i>Radix Scutellaria</i>, <i>Flos Lonicera</i>, and <i>Fructus Forsythiae</i>. The present invention further provides a pharmaceutical composition comprising baicalin, chlorogenic acid and forsythiaside in isolated and purified form. The present invention further provides certain novel derivatives of baicalin, chlorogenic acid and forsythiaside. The present invention additionally provides processes for preparing the extracts as well as baicalin, chlorogenic acid, and forsythiaside in isolated and purified forms.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**PHARMACEUTICAL COMPOSITIONS
AND METHOD OF USING SAME**

TECHNICAL FIELD OF THE INVENTION

5 The present invention relates to pharmaceutical compositions having anitiviral, antibacterial, or immunomodulating property in general, and in particular to pharmaceutical compositions useful in the treatment or prevention of infection by a parainfluenza or respiratory
10 syncytial virus.

BACKGROUND OF THE INVENTION

 The pharmaceutical industry has been developing for many years medicines for the treatment of viral or
15 bacterial infections such as the common cold, flu, and bronchitis. Natural and synthetic products have been considered as possible candidates for these drugs. See, for example, U.S. Patent 4,352,792, which discloses certain 3-alkoxyflavones reportedly useful in treating
20 mammals having infections caused by the human rhinoviruses, enteroviruses and influenzaviruses. U.S. Patent 4,465,673 discloses that 1,2,3,6-pentagalloylglucose is effective against certain viruses, for example, herpes simplex virus. Wang et al. (CJIM,
25 2(3), 162-165 (1996)) states that a certain formulation called Shuanghuanglian (SHL) is useful for the treatment of acute respiratory tract infection. Wang et al. merely states that the "recipe of SHL consists of Flos Lonicerae, Radix Scutellariae and Fructus Fosythiae" and fails to
30 provide a teaching that would allow those skilled in the art to prepare SHL.

 Although there are many drugs, over the counter as well as prescription, available to the public, there is a continuing need for drugs which are effective against
35 these diseases without causing significant side effects.

 These and other objects and advantages of the present invention, as well as additional inventive features, will

be apparent from the description of the invention provided herein.


SUMMARY OF THE INVENTION

5 The foregoing needs have been fulfilled to a great extent by the present invention which provides a pharmaceutical composition comprising an extract or combination of extracts having an antiviral, antibacterial, or immunomodulating property, wherein the
10 extract or the combination of extracts is obtained from a combination of plants, wherein the combination of plants includes at least one plant from each of the genres *Labiatae*, *Caprifoliaceae*, and *Oleaceae*.


The present invention also provides a pharmaceutical
15 composition comprising an extract or a combination of extracts having an antiviral, antibacterial, or immunomodulating property, wherein the extract or combination of extracts is obtained from a combination of *Radix Scutellariae*, *Flos Lonicerae* and *Fructus Forsythiae*.
20 This extract is referred to herein as the RFF extract.

The present invention further provides a pharmaceutical composition having an antiviral, antibacterial, or immunomodulating property, comprising baicalin, chlorogenic acid, and forsythiaside,
25 particularly in isolated and purified form.

The present invention further provides a compound of the formula I or II:



(I)



(II)

wherein R_1 is a sugar residue having 4-6 carbon atoms, optionally having a substituent selected from the group consisting of COOH and an amide, ester, and salt thereof, 5 cyano, amino, substituted amino, alkyl, hydroxyalkyl, halo, and nitro; R_2 and R_3 are selected from the group consisting of hydroxy, alkyloxy, alkyl carbonyl, alkyl carbonyloxy, alkyloxy carbonyl, halo, cyano, nitro, and 10 amino; wherein one or more aromatic hydrogens may be replaced with a substituent selected from the group consisting of hydroxy, cyano, amino, substituted amino, alkyl, hydroxyalkyl, halo, and nitro; and wherein R_4 is a cyclohexyl or disaccharide moiety optionally substituted 15 with one or more substituents selected from the group consisting of COOH and an amide, ester, and salt thereof, cyano, amino, alkyl amino, alkyl, alkyloxy, hydroxyalkyl, hydroxy, halo, nitro, aryl, alkyl aryl, and aryl alkyl wherein said aryl group may be further substituted with 20 one or more substituents selected from the group consisting of COOH and an amide, ester, and salt thereof, cyano, amino, alkyl amino, alkyl, hydroxyalkyl, hydroxy, alkyloxy, halo, and nitro, with the proviso that the compound is not baicalin, chlorogenic acid or 25 foythiaside.

The present invention further provides a pharmaceutical composition comprising a pharmaceutically acceptable carrier and a compound of formula I or II.

The present invention further provides a method of treating or preventing a viral or bacterial infection or enhancing the immune response in a mammal comprising administering to the mammal a pharmaceutical composition
5 of the present invention.

The present invention further provides a process for preparing the RFF extract comprising the steps of:

(1) preparing an extract of *Radix Scutellariae*, which comprises the steps of:

- 10 (a) extracting *Radix Scutellariae* with water at a temperature above about 25°C to provide a first extract;
- (b) concentrating the first extract to provide a concentrate having a relative density of from about
15 1.20 to about 1.25 at 80°C;
- (c) adjusting the pH of the concentrate to a level of from about 1 to about 2 at 80°C;
- (d) cooling the concentrate to produce a precipitate and recovering the precipitate;
- 20 (e) mixing the precipitate with water to obtain a paste;
- (f) adjusting the pH of the paste to about 7;
- (g) dissolving the paste in ethanol to obtain a first ethanolic solution; and optionally filtering the
25 solution to remove any suspended impurities; and
- (h) concentrating the first ethanolic solution to obtain an extract of *Radix scutellariae* having a relative density less than about 1.03 at 40°C; and
- (2) preparing a combined extract of *Flos Lonicerae*
30 and *Fructus Forsythiae* which comprises the steps of:
- (a) extracting a mixture of *Flos lonicerae* and *Fructus forsythiae* with water at a temperature above about 25°C to provide a second extract;

(b) concentrating the second extract to provide a second concentrate having a relative density of from about 1.20 to about 1.25 at 80°C;

(c) cooling the second concentrate to about 40°C;

5 (d) adding ethanol to the second concentrate to obtain a second ethanolic solution and removing any precipitate from the solution;

(e) concentrating the second ethanolic solution to remove the ethanol and recovering the combined
10 extract of *Flos Lonicerae* and *Fructus Forsythiae*; and

(3) combining the extract of *Radix Scutellariae* and the combined extract of *Flos Lonicerae* and *Fructus Forsythiae* to obtain the RFF extract.

The present invention further provides a process for
15 isolating baicalin in a purified form from a plant material containing baicalin comprising:

(a) contacting the plant material with water at an elevated temperature to produce a water extract;

(b) recovering the water extract;

20 (c) adjusting the pH of the water extract to an acidic pH;

(d) diluting the extract with water to obtain a precipitate;

(e) dissolving the precipitate in an aqueous alkali
25 solution to obtain a neutral solution of the precipitate;

(f) diluting the neutral solution with an alcohol to obtain an alcoholic solution;

(g) adjusting the pH of the alcoholic solution to an acidic pH and obtaining a precipitate containing baicalin;

30 and

(h) recovering the purified baicalin.

The present invention further provides a process for isolating chlorogenic acid in a purified form from a plant material containing chlorogenic acid comprising:

(a) contacting the plant material with water to obtain a water extract;

(b) partitioning the water extract with an organic solvent to obtain an organic phase containing chlorogenic acid;

(c) concentrating the organic phase to obtain a concentrate;

(d) purifying the concentrate on a chromatographic column; and

(e) recovering the purified chlorogenic acid.

The present invention further provides a process for isolating forsythiaside in a purified form from a plant material containing forsythiaside comprising:

(a) contacting the plant material with water at an elevated temperature to obtain a water extract;

(b) concentrating the water extract to obtain a concentrate;

(c) contacting the extract with an organic solvent to obtain an organic solution;

(d) recovering and concentrating the organic solution to obtain a residue containing forsythiaside;

(e) purifying the residue on a chromatographic column; and

(f) recovering the purified forsythiaside.

While the invention has been described and disclosed below in connection with certain preferred embodiments and procedures, it is not intended to limit the invention to those specific embodiments. Rather it is intended to cover all such alternative embodiments and modifications as fall within the spirit and scope of the invention.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention provides a pharmaceutical composition comprising an extract or combination of extracts having an antiviral, antibacterial, or

immunomodulating property. The extract or combination of extracts is obtained from a combination of plants wherein the combination of plants includes at least one plant from each of the genres *Labiatae*, *Caprifoliaceae*, and

5 *Oleaceae*.

Any suitable subgenus or species of these plant genres can be used to obtain the extracts or combination of extracts for the inventive pharmaceutical composition. A particular example of a suitable subgenus of *Labiatae* is
10 *Scutellaria* L., a particular example of a suitable subgenus of *Caprifoliaceae* is *Lonicera* L., and a particular example of a suitable subgenus of *Oleaceae* is a *Forsythia* V. Any suitable species can be used. A preferred *Scutellaria* L. specie is *Scutellaria baicalensis*
15 Georgi, a preferred *Lonicera* L. specie is *Lonicerae Japonica* (Thunb.), and a preferred *Forsythia* V. specie is *Forsythia Suspensa* (Thunb.) Vahl. Thus, a preferred extract is obtained from a combination of plants that includes *Scutellaria baicalensis* Georgi, *Lonicera Japonica*
20 (Thunb.), and *Forsythia Suspensa* (Thunb.) Vahl.

Any suitable part of the plant can be used. For example, leaves, twigs, branches, bark, roots, flowers, and fruits can be used.

A description of the above plant species follows.
25 *Scutellaria baicalensis* Georgi is a perennial herbaceous plant with robust and cone-shaped root, quadrigonal stem, 30-60 cm high opposite leaf, terminal panicle, labiate corolla, and purpureous, with a fluorescence time of from July to August and a bearing time of from August to
30 September. This plant grows in the Hebei, Shanxi, and Heilongjiang provinces of the People's Republic of China (PRC). A preferred part of the plant is the root, also known as *Radix Scutellariae*.

Lonicera japonica Thunb. is a semi-evergreen vine
35 with dark red young branch papery leaf, ovate 3-5 cm long

corolla white, yellow late, 3-4.5 cm long, labiate, with a fluorescence time of from April to June and a bearing time of from October to November. This plant grows in the Shandong and Henan provinces of PRC. A preferred part of the plant is the flower, also known as *Flos Lonicera*.

Forsythia Suspensa (Thunb.) Vahl is a deciduous vine with 2-4 m high branch spreading or extended with a little trailing monophyllous, opposite leaf, flower blossoms before leaf shows, axillary about 2.6 cm long capsule narrow ovate 1.5 cm long, with a fluorescence time of from March to May and a bearing time of from July to August. This plant grows in the Shanxi, Henan, and Shandong provinces of PRC. A preferred part of the plant is the fruit, also known as *Fructus Forsythiae*.

An extract containing the combined extracts of *Radix Scutellariae* (R), *Flos Lonicera* (F) and *Fructus Forsythiae* (F), RFF, is particularly preferred. The RFF extract possesses antiviral, antibacterial, and immunostimulating properties.

The present invention further provides a process for preparing an RFF extract. The RFF extract can be prepared by any suitable method. For example, an extract can be prepared from each plant and a combined RFF extract can be prepared by suitably combining the extracts from the other plants. A preferred process is set forth below. The process comprises the steps of:

(1) preparing an extract of *Radix Scutellariae*, which comprises the steps of:

(a) extracting *Radix Scutellariae* with water at a temperature above about 25°C to provide a first extract;

(b) concentrating the first extract to provide a concentrate having a relative density of from about 1.20 to about 1.25 at 80°C;

- (c) adjusting the pH of the concentrate to a level of from about 1 to about 2 at 80°C;
- (d) cooling the concentrate to produce a precipitate and recovering the precipitate;
- 5 (e) mixing the precipitate with water to obtain a paste;
- (f) adjusting the pH of the paste to about 7;
- (g) dissolving the paste in ethanol to obtain a first ethanolic solution; and optionally filtering the
- 10 solution to remove any suspended impurities; and
- (h) concentrating the first ethanolic solution to obtain an extract of *Radix Scutellariae* having a relative density less than about 1.03 at 40°C; and
- (2) preparing a combined extract *Flos Lonicerae* and
- 15 *Fructus Forsythiae* which comprises the steps of:
- (a) extracting a mixture of *Flos Lonicerae* and *Fructus Forsythiae* with water at a temperature above about 25°C to provide an extract;
- (b) concentrating the second extract to provide a
- 20 second concentrate having a relative density of from about 1.20 to about 1.25 at 80°C;
- (c) cooling the second concentrate to about 40°C;
- (d) adding ethanol to the second concentrate to obtain a second ethanolic solution and removing any
- 25 precipitate from the solution;
- (e) concentrating the second ethanolic solution to remove the ethanol and recovering the combined extract of *Flos Lonicerae* and *Fructus Forsythiae* ;
- and
- 30 (3) combining the extract of *Radix Scutellariae* and the combined extract of *Flos Lonicerae* and *Fructus Forsythiae* to obtain the RFF extract.

Any suitable alcohol, preferably ethanol, can be used in the above process.

The RFF extract can have any suitable combination of the three plant parts. For example, the plant parts R:F:F can be in the ratio of about 0.5-1:0.5-1:1-2, preferably about 1:1:2.


5 The present invention further provides a pharmaceutical composition comprising baicalin, chlorogenic acid, and forsythiaside, particularly in isolated and purified form. The present inventor has carried out painstaking and extensive research to identify
10 the active ingredients in the plant extracts. Particularly, it has been discovered by the present inventor that baicalin isolated from the *Scutellaria* plant, *Scutellaria baicalensis* Georgi, is effective as a component of the pharmaceutical composition of the present
15 invention.

Further, it has been discovered by the present inventor that chlorogenic acid is an important ingredient of the plant extract of *Lonicera Japonica* (Thunb.) and is effective as a component of the pharmaceutical composition
20 of the present invention.

In addition, it has been discovered by the present inventor that forsythiaside is an important ingredient of the plant extract of *Forsythia Suspensa* (Thunb.) Vahl and is effective as a component of the pharmaceutical
25 composition of the present invention. The formulas of baicalin, chlorogenic acid, and forsythiaside are set forth herein below:


Baicalin

5,6-Dihydroxy-4-oxo-2-phenyl-4H-t-benzopyran-7-yl- β -D-glucopyranosiduronic acid




Chlorogenic acid

3-(3-(3,4-dihydroxyphenyl)-[-oxo-2-propenyl]oxy)-1,4,5-trihydroxy-, [1S-(1a,3b, 4a, 5a)]-cyclohexanecarboxylic acid



Forsythiaside

(3,4-dihydroxy- β -phenothyl-O- α -L-rhamno-pyranosyl
- (1 \rightarrow 6)-4-O-caffeoyl- β -D-glucopyranoside)



The present invention further provides an antiviral, antibacterial, or immunostimulating composition comprising the active ingredients baicalin (B), chlorogenic acid (C), and forsythiaside (F). This composition is referred to herein as the BCF composition. The BCF composition, by virtue of its greater purity of the active ingredients, is as equally as effective as the RFF composition at lower concentrations of the ingredients. Thus, smaller concentrations of the active ingredients can be used in BCF. Any possible side effect is minimized because of the greater purity of the ingredients and the reduced concentration of the ingredients.

The present invention further provides a process for isolating baicalin in a purified form from a plant material containing baicalin comprising:

- (a) contacting the plant material with water at an elevated temperature to produce a water extract;
- (b) recovering the water extract;
- (c) adjusting the pH of the water extract to an acidic pH;

(d) diluting the extract with water to obtain a precipitate;

(e) dissolving the precipitate in an aqueous alkali solution to obtain a neutral solution of the precipitate;

5 (f) diluting the neutral solution with an alcohol to obtain an alcoholic solution;

(g) adjusting the pH of the alcoholic solution to an acidic pH and obtaining a precipitate containing baicalin; and

10 (h) recovering purified baicalin.

Any suitable organic solvent, preferably an alcohol, more preferably ethanol, can be used in the above process.

Thus, for example baicalin preferably is isolated in purified form from *Radix Scutellariae* as follows: The
15 plant part is cut into small pieces and extracted with water at an elevated temperature, for example, at a temperature of from about 60°C to about 100°C, and preferably at about 90°C. The ratio of water-to-plant weight is selected such that water is in excess,
20 preferably about 8 times that of the plant. The extraction can be repeated to insure high recovery of the active ingredient. The extracts are combined to yield a combined extract, which can be filtered to remove any suspended impurities.

25 The filtered extract is then acidified, for example, with conc. HCl, to a pH of about 2. The acidified extract is diluted with excess water to obtain a precipitate, which is then separated from the supernatant, e.g., by filtration. The filtered precipitate then is mixed with
30 an aqueous alkali solution, e.g., a 20% by wt. NaOH solution and the resulting neutral solution is mixed with an organic solvent, preferably ethanol. The resulting ethanol solution is then acidified, e.g., with HCl to a pH of about 2 to obtain baicalin as a precipitate. The

precipitate is then dried at about 80°C for about 4 hours to obtain baicalin in isolated and purified form.

The present invention further provides a process for isolating chlorogenic acid in a purified form from a plant

5 material containing chlorogenic acid comprising:

(a) contacting the plant material with water to obtain a water extract;

(b) partitioning the water extract with an organic solvent to obtain an organic phase containing chlorogenic
10 acid;

(c) concentrating the organic phase to obtain a concentrate;

(d) purifying the concentrate on a chromatographic column; and

15 (e) recovering chlorogenic acid.

Any suitable organic solvent, preferably an alcohol, and more preferably n-butanol, can be used in the above process.

Thus, for example, chlorogenic acid preferably is
20 isolated in purified form from *Flos Lonicerae* as follows:
The plant part is extracted by stirring with excess water, preferably at a 10-fold excess, for about an hour. The water extract is separated from the residue which is contacted with fresh excess water to yield a second
25 extract. The combined extract is filtered to remove any suspended impurities. The extract is concentrated to reduce the volume by a factor of about 2, and the concentrated extract is then extracted with an organic solvent, e.g., n-butanol. After the solvent removal, the
30 solids subjected to a further purification on a chromatographic column, preferably on a polyamide column. The adsorbed product is eluted using a 30% ethanol solution. The ethanol solvent is then recovered on a HPLC column, e.g., a C₁₈ column. Chlorogenic acid that is
35 retained in the column is eluted using a methanol-water

(2:3 v/v) eluant. The solvent is removed from the eluate, e.g., by freeze drying, to recover purified chlorogenic acid.

The present invention further provides a process for isolating forsythiaside in a purified form from a plant material containing forsythiaside comprising:

(a) contacting the plant material with water at an elevated temperature to obtain a water extract;

(b) concentrating the water extract to obtain a concentrate;

(c) contacting the extract with an organic solvent to obtain an organic solution;

(d) recovering and concentrating the organic solution to obtain a residue containing forsythiaside;

(e) purifying the residue on a chromatographic column; and

(f) recovering purified forsythiaside.

Any suitable organic solvent, preferably an alcohol and more preferably methanol, can be used in the above process.

Thus, for example, forsythiaside preferably is isolated in purified form from *Fructus Forsythiae* as follows: The plant part is extracted with water at an elevated temperature, preferably at about 100°C. The water extract is separated from the residue. The residue is extracted again with water at an elevated temperature, and the extract is collected. The combined extract is filtered to remove any insoluble impurities and then concentrated to reduce the volume. The concentrated extract is extracted with a hot organic solvent, e.g., methanol. The organic phase is separated and the solvent is removed to yield a residue which is then subjected to chromatographic purification on a silica gel column using chloroform-methanol (5:1 v/v) gradient as the eluant. The eluate is concentrated on a HPLC column, e.g., on a C₁₈

column. Forsythiaside adsorbed on the column is eluted using methanol-water (1:1 v/v) to obtain an eluate containing purified forsythiaside. The eluate is concentrated and freeze-dried to obtain forsythiaside in isolated and purified form.

The pharmaceutical compositions of the present invention are particularly suitable for treating or preventing an infection by bacteria and other viruses that affect the respiratory system. Particular examples of these viruses include parainfluenza virus types 1, 2, 3, and 4, and respiratory syncytial virus (RSV) of the long type.


The pharmaceutical compositions of the present invention are particularly suitable for treating or preventing infection by bacteria selected from the group consisting of *Vibrio metchnikovii*, *Edwardsiella tarda*, *Proteus vulgaris*, *Shigalla flexneri*, *Shigalla sonnei*, *Staphylococcus aureus*, *Proteus mirabilis*, *Citrobacter freundii*, Group D streptococcus, *P. shigelloides*, *Aeromonas hydrophilia*, *Salmonella typhi*, *Pseudomonas aeruginosa*, *E. coli*, *H. alvei*, *Enterobacter cloacea*, *Staphylococcus epidermis*, *Streptococcus pneumoniae*, *Branhamella*, and *Bacillus subtilis*.

The pharmaceutical compositions of the present invention are further capable of enhancing the natural killer (NK) cell activity or augmenting the production of interferon-alpha.


The present invention further provides a method of treating or preventing a viral or bacterial infection in a mammal comprising administering to said mammal a pharmaceutical composition as described above.

The present invention further provides a method of enhancing the NK cell activity or augmenting the production of interferon-alpha of a mammal comprising administering to the mammal a pharmaceutical composition described above.

The present invention further provides a compound of the formula I or II:



(I)



(II)

- 5 wherein R_1 is a sugar residue having 4-6 carbon atoms optionally having a substituent selected from the group consisting of COOH and an amide, ester, and salt thereof, cyano, amino, substituted amino, alkyl, hydroxyalkyl,
- 10 halo, and nitro; R_2 and R_3 are selected from the group consisting of hydroxy, alkyloxy, alkyl carbonyl, alkyl carbonyloxy, alkyloxy carbonyl, halo, cyano, nitro, and amino; wherein one or more aromatic hydrogens may be replaced with a substituent selected from the group
- 15 consisting of hydroxy, cyano, amino, substituted amino, alkyl, hydroxyalkyl, halo, and nitro; and wherein R_4 is a cyclohexyl or disaccharide moiety, optionally substituted with one or more substituents selected from the group consisting of COOH and an amide, ester, and salt thereof,
- 20 cyano, amino, alkyl amino, alkyl, alkyloxy, hydroxyalkyl, hydroxy, halo, nitro, aryl, alkyl aryl, and aryl alkyl, wherein said aryl group may be further substituted with one or more substituents selected from the group consisting of COOH and an amide, ester, and salt thereof,
- 25 cyano, amino, alkyl amino, alkyl, hydroxyalkyl, hydroxy,

alkyloxy, halo, and nitro, with the proviso that the compound is not baicalin, chlorogenic acid or foythiaside.

Preferred examples of the above compounds include those wherein R_2 and R_3 are hydroxy. Preferred examples of the compounds include those wherein aryl alkyl is phenylethyl. Preferred examples of the compounds of the present invention include those wherein R_4 is cyclohexyl substituted with 1-3 hydroxyl groups and a carboxyl. A further preferred R_4 is a cyclohexyl substituted with 3 hydroxyl groups and a carboxyl. Other preferred examples of the compounds of the present invention include those wherein R_4 is a disaccharide, particularly a rhamnopyranosyl glucopyranoside. A further preferred disaccharide is 3,4-dihydroxy- β -phenothyl-O-L-rhamnopyranosyl-(1 \rightarrow 6)- β -D-glucopyranoside.

The term "sugar" herein refers to any suitable sugar moiety known to those of ordinary skill in the art including mono and disaccharide moieties. Preferred sugar moieties include monosaccharides having 4-6 carbon atoms. Hexoses are particularly preferred. Glucose is a preferred hexose. A glucose moiety having a carboxyl is further preferred. A preferred R_1 is β -D-glucopyranosiduronic acid.

The term "alkyl" herein refers to an alkyl moiety of any suitable chain length. Preferred alkyl moieties include those having 1-6 carbon atoms. The term "aryl" herein refers to aromatic groups having 1-3 rings, preferably a phenyl.

The compounds of formula I and II can be prepared by methods known to those of ordinary skill in the art. For example, these compounds can be prepared by suitable modification of baicalin, chlorogenic acid, or foythiaside. Thus, for example, a hydroxyl group can be converted to an ether by converting the hydroxyl to

alkoxide followed by the reaction of the alkoxide with an alkyl iodide. The aromatic ring hydrogens can be replaced by substituents such as nitro, e.g., by an electrophilic substitution reaction. An amino group can be obtained by the reduction of the nitro group, e.g., by platinum/hydrogen. The aromatic ring positions can be halogenated, e.g., photochemically. An alkyl group can be introduced into the aromatic ring via Friedel-Craft alkylation procedures known to those of ordinary skill in the art. A carboxyl group can be introduced in the sugar moiety by oxidation, e.g., of the hydroxymethyl group. A carboxyl group can be converted to the amide by reaction with an amine, followed by dehydration. A carboxyl group can be converted to the ester by reaction with an alcohol under acidic conditions. Other derivatives can also be prepared by methods known to those of ordinary skill in the art including those disclosed in common text books; see, e.g., Advanced Organic Chemistry, J. March, 3^d Ed. (1985).

The present invention further provides a pharmaceutical composition comprising a pharmaceutically acceptable carrier and an effective (e.g., therapeutically or prophylactically effective) amount of at least one of the compounds set forth above.

The present invention further provides a method of treating or preventing a viral or bacterial infection in a mammal comprising administering to the mammal an effective amount (e.g., therapeutically or prophylactically effective) of at least one of the compounds set forth above.

The present invention further provides a method of enhancing the NK cell activity or augmenting the production of interferon-alpha of a mammal comprising administering to the mammal at least one of the compounds set forth above.

The pharmaceutically acceptable carriers described herein, for example, vehicles, adjuvants, excipients, or diluents, are well-known to those who are skilled in the art and are readily available to the public. It is preferred
5 that the pharmaceutically acceptable carrier be one which is chemically inert to the active compounds and one which has no detrimental side effects or toxicity under the conditions of use.

The choice of carrier will be determined in part by the
10 particular active agent, as well as by the particular method used to administer the composition. Accordingly, there is a wide variety of suitable formulations of the pharmaceutical composition of the present invention. The following
15 formulations for oral, aerosol, parenteral, subcutaneous, intravenous, intraarterial, intramuscular, interperitoneal, intrathecal, rectal, and vaginal administration are merely exemplary and are in no way limiting.

Formulations suitable for oral administration can consist of (a) liquid solutions, such as an effective amount
20 of the compound dissolved in diluents, such as water, saline, or orange juice; (b) capsules, sachets, tablets, lozenges, and troches, each containing a predetermined amount of the active ingredient, as solids or granules; (c) powders; (d) suspensions in an appropriate liquid; and (e)
25 suitable emulsions. Liquid formulations may include diluents, such as water and alcohols, for example, ethanol, benzyl alcohol, and the polyethylene alcohols, either with or without the addition of a pharmaceutically acceptable surfactant, suspending agent, or emulsifying agent. Capsule
30 forms can be of the ordinary hard- or soft-shelled gelatin type containing, for example, surfactants, lubricants, and inert fillers, such as lactose, sucrose, calcium phosphate, and corn starch. Tablet forms can include one or more of
35 alginic acid, microcrystalline cellulose, acacia, gelatin, guar gum, colloidal silicon dioxide, croscarmellose sodium,

5 talc, magnesium stearate, calcium stearate, zinc stearate,
stearic acid, and other excipients, colorants, diluents,
buffering agents, disintegrating agents, moistening agents,
preservatives, flavoring agents, and pharmacologically
compatible carriers. Lozenge forms can comprise the active
ingredient in a flavor, usually sucrose and acacia or
tragacanth, as well as pastilles comprising the active
ingredient in an inert base, such as gelatin and glycerin,
or sucrose and acacia, emulsions, gels, and the like
10 containing, in addition to the active ingredient, such
carriers as are known in the art.

The compounds of the present invention, alone or in
combination with other suitable components, can be made into
aerosol formulations to be administered via inhalation.
15 These aerosol formulations can be placed into pressurized
acceptable propellants, such as dichlorodifluoromethane,
propane, nitrogen, and the like. They also may be formulated
as pharmaceuticals for non-pressured preparations, such as
in a nebulizer or an atomizer.

20 Formulations suitable for parenteral administration
include aqueous and non-aqueous, isotonic sterile injection
solutions, which can contain anti-oxidants, buffers,
bacteriostats, and solutes that render the formulation
isotonic with the blood of the intended recipient, and
25 aqueous and non-aqueous sterile suspensions that can include
suspending agents, solubilizers, thickening agents,
stabilizers, and preservatives. The compound can be
administered in a physiologically acceptable diluent in a
pharmaceutical carrier, such as a sterile liquid or mixture
30 of liquids, including water, saline, aqueous dextrose and
related sugar solutions, an alcohol, such as ethanol,
isopropanol, or hexadecyl alcohol, glycols, such as
propylene glycol or polyethylene glycol, glycerol ketals,
such as 2,2-dimethyl-1,3-dioxolane-4-methanol, ethers, such
35 as poly(ethyleneglycol) 400, an oil, a fatty acid, a fatty
acid ester or glyceride, or an acetylated fatty acid

glyceride with or without the addition of a pharmaceutically acceptable surfactant, such as a soap or a detergent, suspending agent, such as pectin, carbomers, methylcellulose, hydroxypropylmethylcellulose, or carboxymethylcellulose, or emulsifying agents and other pharmaceutical adjuvants.

Oils, which can be used in parenteral formulations include petroleum, animal, vegetable, or synthetic oils. Specific examples of oils include peanut, soybean, sesame, cottonseed, corn, olive, petrolatum, and mineral. Suitable fatty acids for use in parenteral formulations include oleic acid, stearic acid, and isostearic acid. Ethyl oleate and isopropyl myristate are examples of suitable fatty acid esters. Suitable soaps for use in parenteral formulations include fatty alkali metal, ammonium, and triethanolamine salts, and suitable detergents include (a) cationic detergents such as, for example, dimethyl dialkyl ammonium halides, and alkyl pyridinium halides, (b) anionic detergents such as, for example, alkyl, aryl, and olefin sulfonates, alkyl, olefin, ether, and monoglyceride sulfates, and sulfosuccinates, (c) nonionic detergents such as, for example, fatty amine oxides, fatty acid alkanolamides, and polyoxyethylenepolypropylene copolymers, (d) amphoteric detergents such as, for example, alkyl-b-aminopropionates, and 2-alkyl-imidazoline quaternary ammonium salts, and (e) mixtures thereof.

The parenteral formulations will typically contain from about 0.5 to about 25% by weight of the active ingredient in solution. Suitable preservatives and buffers can be used in such formulations. In order to minimize or eliminate irritation at the site of injection, such compositions may contain one or more nonionic surfactants. The quantity of surfactant in such formulations ranges from about 5 to about 15% by weight. Suitable surfactants include polyethylene sorbitan fatty acid esters, such as sorbitan monooleate and the high molecular weight adducts of ethylene oxide with a

hydrophobic base, formed by the condensation of propylene oxide with propylene glycol. The parenteral formulations can be presented in unit-dose or multi-dose sealed containers, such as ampoules and vials, and can be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example, water, for injections, immediately prior to use. Extemporaneous injection solutions and suspensions can be prepared from sterile powders, granules, and tablets of the kind previously described.

The compounds of the present invention may be made into injectable formulations. The requirements for effective pharmaceutical carriers for injectable compositions are well known to those of ordinary skill in the art. See, Pharmaceutics and Pharmacy Practice, J.B. Lippincott Co., Philadelphia, PA, Banker and Chalmers, eds., pages 238-250 (1982), and ASHP Handbook on Injectable Drugs, Toissel, 4th ed., pages 622-630 (1986).

Additionally, the compounds of the present invention may be made into suppositories by mixing with a variety of bases, such as emulsifying bases or water-soluble bases. Formulations suitable for vaginal administration may be presented as pessaries, tampons, creams, gels, pastes, foams, or spray formulas containing, in addition to the active ingredient, such carriers as are known in the art to be appropriate.

Suitable doses and dosage regimens can be determined by conventional range-finding techniques known to those of ordinary skill in the art. Generally, treatment is initiated with smaller dosages, which are less than the optimum dose of the compound. Thereafter, the dosage is increased by small increments until the optimum effect under the circumstances is reached. For convenience, the total daily dosage may be divided and administered in portions during the day if desired. In proper doses and with suitable administration of certain compounds, the present

invention provides for a wide range of responses. The dosages range from about 0.001 to about 1000 mg/kg body weight of the animal being treated/day. Preferred dosages range from about 0.01 to about 10 mg/kg body weight/day, and
5 further preferred dosages range from about 0.01 to about 1 mg/kg body weight/day.

The following examples further illustrate the present invention but, of course, should not be construed as in any way limiting its scope.

10

EXAMPLE 1

This example illustrates the preparation of the RFF extract. The RFF extract was prepared by combining the extract of *Radix Scutellariae* and the combined extract of
15 *Flos Lonicera* and *Fructus Fosythiae*. The materials used and the method of preparation are set forth below.

The extract of *Radix Scutellariae* was prepared as follows:

The plant part (500 g) was cut into small slices, and
20 sufficient water was added to cover the slices. The combination was heated, and the water was allowed to boil for 2 hours. The water extract was separated and saved. Sufficient amount of water again was added to cover the residue slices, and the water was allowed to boil for 1
25 hour. The second extract also was saved. The residue was extracted with boiling water once more to obtain a third extract.

The three extracts were combined and concentrated by boiling until the relative density of the liquid was in
30 the range of 1.20-1.25 at 80°C. The pH of the resulting liquid was adjusted to 1.0-2.0 with HCl (2 mol/L) at 80°C. The liquid was stored at 80°C for 1 hour and then allowed to sit at room temperature for 24 hours. The supernatant solution was discarded, and a small amount of water was
35 added to the residue which was then beaten to form a

paste. The pH of the paste was adjusted to 7.0 with 40% NaOH, and an equal volume of 95% ethanol was added to the mixture. The mixture was stirred until the reconstituted residue dissolved. The resulting solution was filtered.

- 5 The solution was then concentrated to have a relative density of no less than 1.03 at 40°C. This solution was allowed to cool to room temperature. This is the extract of *Radix Scutellariae*.

- The combined extract of *Flos Lonicerae* and *Fructus Forsythiae* was prepared as follows:

- 10 *Flos Lonicerae* (500 g) and *Fructus Forsythiae* (1000 g) were bathed in warm water for 30 minutes, and then the bath water was boiled for 1.5 hours. The water extract was separated and saved. The residue was boiled again in
15 water for 1.5 hours. The extract was saved. The two extracts were combined and concentrated by boiling to attain a solution of relative density of 1.20-1.25 at 80°C. The solution was cooled to 40°C, and ethanol was added gradually and with stirring to attain a concentration of
20 75% ethanol. The mixture was allowed to stand for 24 hours and then filtered to remove the precipitate. The precipitate was rinsed with 75% ethanol and filtered. All the filtered liquids were combined, and the ethanol was removed until the residual liquid was free of ethanol
25 odor. The residual liquid was added to the precipitate again with ethanol to attain a concentration of 85% ethanol. The mixture was allowed to stand for 24 hours and filtered. The precipitate was rinsed with 85% ethanol and filtered. All the filtered liquids were combined, and
30 the ethanol was recovered from the filtered liquid until ethanol odor could not be detected in the extract.

The extract of *Radix Scutellariae* prepared earlier was combined with the extract prepared above, and the combination was concentrated by boiling to 2000 ml.

EXAMPLE 2

This example illustrates the virostatic and bacteriostatic activity as well as the immunomodulating effects of the pharmaceutical compositions of the present invention. One embodiment of the present invention, RFF, is a composition comprising an extract of a mixture of *Radix Scutellariae*, *Flos Lonicerae*, *Fructus Forsythiae*. Another embodiment of the present invention is BCF which is a mixture of the active ingredients of the above three plants, i.e., a mixture of baicalin, chlorogenic acid and forsythiaside. The materials used, the method of testing, and the results obtained are discussed below:

Materials and Methods

1. Pharmaceutical Compositions
- 15 RFF - each ml contained an extract from a mixture containing 0.25 g *Radix scutellariae*, 0.25 g *Flos lonicerae* and 0.5 g *Fructus forsythiae*.
BCF - each ml contained 2 mg baicalin, 1 mg chlorogenic acid and 1 mg forsythiaside.
- 20 Cell maintenance media were used as control.

2. Cell types

K562, Wsh, Hela, and Hep-2 cells were employed in this study. The cells were maintained and passaged under conditions known to those skilled in the art.

5

3. Viruses

The following viruses were tested: parainfluenza virus types: 1, 2, 3, and 4; and respiratory syncytial virus of the long type.

10

These viruses were obtained from the China Institute of Medical Sciences Laboratory and were maintained according to established protocols known to those skilled in the art. Their TCD50 and agglutination titers were determined by published methods.

15

4. Bacterial strains

The bacterial strains tested are set forth in Table 1. These strains were obtained from the China Institute of Medical Sciences and maintained according to established protocols.

20

5. NK cell activity assay

To each well of the 96-well plate, 0.1 ml effector cells (containing 5000 lymphocytes) and 0.1 ml target cells (containing 500 K562 cells) were added. Each sample was made in triplicate. The cells were incubated at 37°C, 5% CO₂ for 24 hours. The serum was then removed, following which 0.1 ml DNAase (100 mg/ml) and 0.1 ml trypsin (4 mg/ml) were added, and the cells were further incubated for another 24 hours. The cells were then pulsed with ³H-thymidine for 4 hours and harvested onto fiber filters, and the amount of ³H-thymidine (CPM) on the filter was determined by scintillation counting. The wells that did not have added target cells served as control.

30

35

HTdR release % = Target cell (CPM) - control (CPM)

6. Determination of the activity of Inteferon-alpha

The tested specimens were serially diluted in duplicate and added to a monolayer of Wsh cells. Normal
5 cells with and without added virus were used as control cells. The specimens were incubated at 37°C for 24 hours, and then 200TCD50VSV were added. The cells with added virus were then scored when they showed +++ virus induced pathological changes. The results obtained are discussed
10 below.

Results

1. Virostatic studies

Both BCF and RFF were diluted to 0.5% by wt.
15 concentration with cell maintenance media. Three groups (Groups A-C) of tests were performed.
Group A: The above solutions were added individually to monolayer cells, and the cells were incubated for 2 hours at 37°C. 100TCD50 virus suspensions were added and then
20 removed and replaced with cell maintenance media.
Group B: The above solutions were individually mixed with 100TCD50 virus suspensions incubated at 37°C for 1 hour and then added to monolayer cells. The cells were kept at 37°C for 1 hour and then replaced with cell maintenance media.
25 Group C: After the monolayer cells were exposed to the virus for 24 hours, of either BCF or RFF were added to the cells.

The controls were normal cells exposed to the virus and not to BCF or RFF. All specimens were incubated at
30 37°C for 7 days, and then the virus was titered. The compositions were found to have antiviral activity if the viral load between the treatment groups and the control groups was reduced four-fold. The results obtained are set forth in Table 1.

35

Table 1: Antiviral activity of RFF and BCF

Virus	BCF			RFF			CONTROL		
	A	B	C	A	B	C	A	B	C
Parainfl. 1	-	+	+	-	+	+	-	-	-
Parainfl. 2	-	+	+	-	+	+	-	-	-
Parainfl. 3	-	+	+	-	+	+	-	-	-
Parainfl. 4	-	+	+	-	+	+	-	-	-
RSV	-	+	+	-	+	+	-	-	-

5 '+' means an antiviral effect was observed, "-" means no antiviral effect was observed. "Parainfl." stands for parainfluenza virus.

10 The results were obtained in triplicate and were unequivocal. The results demonstrated that RFF and BCF were effective in reducing the virotoxic effects of parainfluenza virus types 1-4 as well as RSV when RFF and BCF were administered with the virus (Group B) or after viral exposure (Group C).

2. Bacteriostatic studies

A. The bacteria were allowed to grow for 16-24 hours and then the bactererial density was determined by established methods with the final bacterial concentration kept at one billion/ml. The bacteria grew in both liquid and solid media, and bacterial densities were measured after 16-24 hours. The results obtained are set forth in Table 2.

Table 2: Antibacterial effect of RFF and BCF

Bacteria	BCF ¹				RFF ¹				Control ¹			
	Concentration (%)				Concentration (%)				Concentration (%)			
	20	10	5	2.5	20	10	5	2.5	20	10	5	2.5
<i>Serratia marcescens</i>	+	+	+	+	+	+	+	+	+	+	+	+
<i>Vibrio metchnikovli</i>	-	-	-	-	-	-	-	+	+	+	+	+
<i>Edwardsiella tarda</i>	-	-	-	-	-	-	-	+	+	+	+	+
<i>Proteus vulgaris</i>	-	-	-	+	-	-	-	+	+	+	+	+
<i>Shigalla flexneri</i>	-	-	+	+	-	-	+	+	+	+	+	+
<i>Staphylococcus aureus</i>	-	-	-	-	-	-	-	-	+	+	+	+
<i>Proteus mirabilis</i>	-	-	-	+	-	-	-	+	+	+	+	+
<i>Citrobacter freundil</i>	-	-	+	+	-	-	+	+	+	+	+	+
Group D streptococcus	-	+	+	+	-	+	+	+	+	+	+	+
<i>P. shigelloides</i>	-	-	-	-	-	-	-	-	+	+	+	+
<i>Aeromonas hydrophilia</i>	-	-	-	+	-	-	-	+	+	+	+	+
<i>Salmonella typhi</i>	-	-	+	+	-	-	-	+	+	+	+	+
<i>Pseudomonas aeruginosa</i>	-	-	+	+	-	-	+	+	+	+	+	+
<i>Escherichia Coli</i>	-	+	+	+	-	+	+	+	+	+	+	+

Bacteria	BCF ¹				RFF ¹				Control ¹			
	Concentration (%)				Concentration (%)				Concentration (%)			
	20	10	5	2.5	20	10	5	2.5	20	10	5	2.5
<i>H. alvei</i>	-	+	+	+	-	+	+	+	+	+	+	+
<i>Enterobacter cloacae</i>	-	-	+	+	-	-	+	+	+	+	+	+
<i>Staphylococcus epidermidis</i>	-	-	+	-	-	-	+	+	+	+	+	+
<i>Staphylococcus citreus</i>	+	+	+	+	+	+	+	+	+	+	+	+
<i>Streptococcus pneumoniae</i>	-	-	+	+	-	-	+	+	+	+	+	+
<i>Enterobacter aerogenes</i>	+	+	+	+	+	+	+	+	+	+	+	+
<i>Branhamella</i>	-	-	+	+	-	-	+	+	+	+	+	+
<i>Bacillus subtilis</i>	-	-	+	+	-	-	+	+	+	+	+	+

¹ - indicates no bacterial growth was observed.

¹ + indicates bacterial growth was observed.

- 1 The BCF, RFF or control solution was prepared by diluting the original solutions (see Materials and Methods - 1. Pharmaceutical compositions) by a factor of 5, 10, 20, or 40 to obtain 20, 10, 5, or 2.5% BCF, RFF, or control solution, respectively.

The above results demonstrate that each of BCF and RFF had bacteriostatic effect on a number of different bacteria. The effect of each of BCF and RFF correlated with the concentration being used, with the least effect
5 seen at the lowest concentration.

3. Anti-Staph. aureus animal studies

A. Staphylococcus aureus was obtained from the China Institute of Medical Sciences Laboratory. The
10 bacteria were cultured in both blood agar and glucose based media. The MCD towards white mice was determined to be 50 million/animal.

B. Animal experiments were carried out on laboratory bred white mice of average weight 18-22 g. The
15 male and female mice were equal in numbers.

C. Anti-Staphylococcus aureus studies

The pharmaceutical compositions used were BCF or RFF as described above. Concentrations used were as follows:
20 $2 \mu\text{g}/0.1 \text{ ml}$, $4 \mu\text{g}/0.1 \text{ ml}$, $6 \mu\text{g}/0.1 \text{ ml}$, $8 \mu\text{g}/0.1 \text{ ml}$., $10 \mu\text{g}/0.1 \text{ ml}$., $12 \mu\text{g}/0.1 \text{ ml}$, and $14 \mu\text{g}/0.1 \text{ ml}$.

The mice (total $n=160$) were divided into a BCF group ($n=70$), RFF group ($n=70$), and control groups ($n=20$). There were two control groups with 10 animals serving as positive and negative controls.

25 Each animal in the BCF group was injected with 50 millions staphylococcus aureus via the tail vein, and the animals were subdivided into seven groups. Each group received a different concentration of BCF, and the medications were given intraperitoneally daily starting 6
30 hours post staphylococcus aureus injections. Seven days later, the survival rate of the animals was determined. Studies using RFF were performed similarly.

The positive control group consisted of 10 animals receiving the bacteria and normal saline, and the negative
35 control group consisted of 10 animals receiving neither bacteria nor a pharmaceutical composition. The ED50 was

determined by the Cou's method. The results obtained are set forth in Tables 3-4.

- 5 Table 3: Results of Anti-Staphylococcus aureus studies with BCF

group	dose ($\mu\text{g}/\text{animal}$)	BCF			
		n	death	mortality rate	logarithm of dose
1	2	10	10	1.0	0.3010
2	4	10	8	0.8	0.6021
3	6	10	6	0.6	0.7782
4	8	10	4	0.4	0.9031
5	10	10	3	0.3	1.0
6	12	10	2	0.2	1.0792
7	14	10	0	0	1.1461

The ED50 was calculated to be 6.6621 $\mu\text{g}/\text{animal}$,
with 95% confidence level between 5.3215 and 8.5901

10 $\mu\text{g}/\text{animal}$.

Table 4: Results of Anti-Staphylococcus aureus studies
with RFF

group	dose ($\mu\text{g}/\text{animal}$)	RFF			
		n	death	mortality rate	logarithm of dose
1	2	10	10	1.0	0.301
2	4	10	8	0.8	0.602
3	6	10	6	0.6	0.778
4	8	10	5	0.5	0.903
5	10	10	3	0.3	1.0
6	12	10	2	0.2	1.0792
7	14	10	0	0	1.1461

The ED50 was calculated to be 6.747 $\mu\text{g}/\text{animal}$, with a
5 95% confidence level between 5.2264 and 8.7136 $\mu\text{g}/\text{animal}$.

The positive control group animals receiving only the
normal saline all died. The negative control group
animals were all alive. The above data demonstrated that
both BCF and RFF had antistaphylococcus aureus activity.

10

4. Effect on peripheral blood NK cell activity

Peripheral blood lymphocytes were obtained from human
volunteers and then divided and prepared as follows:

- Group A: Lymphocytes + 1640 growth media
- 15 Group B: Lymphocytes + 1% RFF 1640 growth media
- Group C: Lymphocytes + 0.5% RFF 1640 growth media
- Group D: Lymphocytes + 0.25% RFF 1640 growth media

The cells were then incubated at 37°C for 48 hours and
the NK cell activity was determined. The results obtained
20 are set forth in Table 5.

25

Table 5: Effect of RFF concentration on NK cell activity

Group	n	NK activity (X +/- SD)	P value
A	20	46.28 +/- 7.6	-
B	20	66.47 +/- 8.36	<0.01
C	20	68.11 +/- 9.6	<0.01
D	20	60.87 +/- 8.2	<0.01

X = mean; SD = standard deviation

- 5 Similar studies were performed using BCF and the results obtained were as follows:

Table 6: Effect of BCF concentration on NK cell activity

Group	n	NK activity (X +/- SD)	P values
A	20	47.28 +/- 8.0	-
B	20	68.48 +/- 7.06	<0.01
C	20	68.44 +/- 9.31	<0.01
D	20	69.81 +/- 7.5	<0.01

10 X = mean; SD = standard deviation

The above results demonstrated that 0.24%-1% RFF or BCF was able to enhance NK cell activity.

- 15 5. Effect on peripheral blood lymphocytes on the production of interferon-alpha

Peripheral blood lymphocytes were exposed to varying concentrations of RFF or BCF in a growth medium and Newcastle disease virus (NDV). The control lymphocytes were exposed to NDV and growth medium and not to RFF or BCF.

- Group A: Peripheral blood lymphocytes + 1640 growth media + NDV (virus added control)
- 25 Group B: Peripheral blood lymphocytes + 1% RFF 1640 growth media + NDV

Group C: Peripheral blood lymphocytes + 0.5% RFF 1640
growth media + NDV

Group D: Peripheral blood lymphocytes + 0.25% RFF 1640
growth media + NDV

5 The results obtained are set forth in Table 7.

Table 7: Effect of RFF on peripheral blood lymphocytes
production of interferon-alpha

10

Group	n	Interferon- alpha (u/ml, X +/- SD)	T
A	20	4182 +/- 272	-
B	20	7260 +/- 804	<0.01
C	20	8844 +/- 912	<0.01
D	20	8680 +/- 863	<0.01

X = mean; SD = standard deviation; u/ml = units/ml

The above results demonstrated that concentrations of
0.25%-1% RFF enhanced the production of interferon-alpha
by peripheral blood lymphocytes.

15 Studies were performed with BCF in a similar manner,
and the results obtained are set forth in Table 8.

Table 8: Effect of BCF on the production of interferon-
alpha by peripheral blood lymphocytes

Group	n	Interferon-alpha (u/ml, X +/- SD)	t
A	20	4182 +/- 272	-
B	20	7360 +/- 814	<0.01
C	20	8744 +/- 913	<0.01
D	20	8686 +/- 843	<0.01

20 X = mean; SD = standard deviation; u/ml = units/ml

The above results demonstrate that BCF in
concentrations ranging from 0.25%-1% enhanced the
production of interferon-alpha by peripheral blood
25 lymphocytes.

6. Effect on NK cell activity and production of interferon - alpha in rabbits

5 Rabbits (total n=60) were divided into a BCF group (n=20), RFF group (n=20) and control group (n=20). In both the BCF and RFF groups, 5 ml of each medication was injected via the neck vein into the rabbits while the control group received 5 ml of normal saline injection. The injections were continued daily for a total of 5 days, and peripheral blood samples were obtained from the animal's neck artery. NK cell activity as well as interferon-alpha concentrations were determined. The results obtained are set forth in Tables 9-10.

10

Table 9: Effect of RFF on NK cell activity and level of interferon - alpha in rabbits

Group	n	NK cell activity X +/- SD	Inteferon (u/ml, X +/- SD)
RFF	20	60.24 +/- 12.13	1536 +/- 134
Control	20	38.92 +/- 8.2	724 +/- 92
t		<0.01	<0.01

5 X = mean; SD = standard deviation; u/ml = units/ml

Table 10: Effect of BCF of NK cell activity and level of interferon-alpha in rabbits

Group	n	NK cell activity X +/- SD	Inteferon, u/ml, X +/- SD
BCF	20	61.15 +/- 11.14	1632 +/- 125
Control	20	38.92 +/- 8.2	724 +/- 92
t		<0.01	<0.01

10 X = mean; SD = standard deviation; u/ml = units/ml

The above results demonstrate that both BCF and RFF enhanced the activity of NK cell activity as well as the production of interferon-alpha by peripheral blood lymphocytes in rabbits.

15 The foregoing confirms that RFF and BCF have virostatic effects towards RSV and Parainfluenza virus types 1-4. Both in vitro and in vivo data support that both RFF and BCF have bacteriostatic effects on staphylococcus aureus. In vitro data also show that RFF
20 and BCF were effective against pathogens such as Proteus, Shigalla, Salmonella, Pseudomonas, Enterobacter cloacae, Streptococcus. Both BCF and RFF enhance NK cell activity as well as augment the production of interferon-alpha by peripheral blood lymphocytes. Interferon-alpha is an
25 important cytokine in resisting viral infection. Interferon-alpha also possesses an immunomodulating function. It enhances the activity of NK cell activity, augments the macrophage-like function and directly

participates in regulating several biomolecules within the cells. NK cells serve an important immunosurveillance function. They participate directly in eliminating viral infected cells as well as the viral particles itself. NK
5 cells are also important in producing interferon-alpha. These two effects help form an effective immunobARRIER in resisting viral related illnesses.

All of the patents and publications cited herein are incorporated in their entirety by reference.

10 While this invention has been described with an emphasis upon the preferred embodiment, it will be obvious to those of ordinary skill in the art that variations of the preferred embodiment may be used and that it is intended that the invention may be practiced otherwise
15 than as specifically described herein. Accordingly, this invention includes all modifications encompassed within the spirit and scope of the invention as defined by the following claims.

WHAT IS CLAIMED IS:

1. A pharmaceutical composition comprising an extract or combination of extracts having an antiviral, antibacterial, or immunomodulating property, wherein said
5 extract or combination of extracts is obtained from a combination of plants which includes at least one plant from each of the genres *Labiatae*, *Caprifoliaceae*, and *Oleaceae*.
- 10 2. The pharmaceutical composition of claim 1, wherein said *Labiatae* is a *Scutellaria L.*, said *Caprifoliaceae* is a *Lonicera L.*, and said *Oleacea* is a *Forsythia V.*
- 15 3. The pharmaceutical composition of claim 2, wherein said *Scutellaria L.* is *Scutellaria baicalensis* Georgi, said *Lonicera L.* is *Lonicerae Japonica* (Thunb.), and said *Forsythia V.* is *Forsythia Suspensa* (Thunb.) Vahl.
- 20 4. The pharmaceutical composition of claim 3, wherein said extract or combination of extracts is obtained from *Radix Scutellaria*, *Flos Lonicera*, and *Fructus Forsythiae*.
- 25 5. The pharmaceutical composition of claim 4, wherein said extract or combination of extracts has antiviral activity against a virus selected from the group consisting of parainfluenza virus types 1, 2, 3, and 4, and respiratory syncytial virus of the long type.
- 30 6. The pharmaceutical composition of claim 4, wherein said extract or combination of extracts has antibacterial activity against a bacteria selected from the group consisting of *Vibrio metchnikovii*, *Edwardsiella*
35 *tarda*, *Proteus vulgaris*, *Shigalla flexneri*, *Shigalla sonnei*, *Staphylococcus aureus*, *Proteus mirabilis*,

Citrobacter freundii, Group D streptococcus, P. shigelloides, Aeromonas hydrophila, Salmonella typhi, Pseudomonas aeruginosa, E. coli, H. alvei, Enterobacter cloacae, Staphylococcus epidermis, Streptococcus pneumoniae, Branhamella, and Bacillus subtilis.

7. The pharmaceutical composition of claim 4, wherein said extract or combination of extracts is capable of enhancing the NK cell activity or augmenting the production of interferon-alpha.

8. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and baicalin, chlorogenic acid, and forsythiaside in isolated and purified form.

9. A method of treating or preventing a viral or bacterial infection in a mammal comprising administering to said mammal a pharmaceutical composition according to claim 1.

10. A method of treating or preventing a viral or bacterial infection in a mammal comprising administering to said mammal a pharmaceutical composition according to claim 4.

11. A method of treating or preventing a viral or bacterial infection in a mammal comprising administering to said mammal a pharmaceutical composition according to claim 8.

12. A method of enhancing the NK cell activity or augmenting the production of interferon-alpha of a mammal comprising administering to said mammal a pharmaceutical composition according to claim 1.


13. A method of enhancing the NK cell activity or augmenting the production of interferon-alpha of a mammal comprising administering to said mammal a pharmaceutical composition according to claim 4.

5


14. A method of enhancing the NK cell activity or augmenting the production of interferon-alpha of a mammal comprising administering to said mammal a pharmaceutical composition according to claim 8.

10

15. A compound of the formula I or II:



(I)



(II)

15

wherein R₁ is a sugar residue having 4-6 carbon atoms optionally having a substituent selected from the group consisting of COOH and an amide, ester, and salt thereof, cyano, amino, substituted amino, alkyl, hydroxyalkyl, halo, and nitro; R₂ and R₃ are selected from the group consisting of hydroxy, alkyloxy, alkyl carbonyl, alkyl carbonyloxy, alkyloxy carbonyl, halo, cyano, nitro, and amino; wherein one or more aromatic hydrogens may be replaced with a substituent selected from the group consisting of hydroxy, cyano, amino, substituted amino,

20

25

alkyl, hydroxyalkyl, halo, and nitro; and wherein R₄ is a cyclohexyl or disaccharide moiety optionally substituted with one or more substituents selected from the group consisting of COOH and an amide, ester, and salt thereof, cyano, amino, alkyl amino, alkyl, alkyloxy, hydroxyalkyl, hydroxy, halo, nitro, aryl, alkyl aryl, and aryl alkyl wherein said aryl group may be further substituted with one or more substituents selected from the group consisting of COOH and an amide, ester, and salt thereof, cyano, amino, alkyl amino, alkyl, hydroxyalkyl, hydroxy, alkyloxy, halo, and nitro, with the proviso that the compound is not baicalin, chlorogenic acid or foythiaside.

15 16. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and an effective amount of at least one compound of claim 15.

20 17. A method of treating or preventing a viral or bacterial infection in a mammal comprising administering to said mammal an effective amount of at least one compound of claim 15.

25 18. A method of enhancing the NK cell activity or augmenting the production of interferon-alpha of a mammal comprising administering to said mammal an effective amount of at least one compound of claim 15.

30 19. A process for preparing an RFF extract from *Radix Scutellariae*, *Flos Lonicerae*, and *Fructus Forsythiae*, said extract having antiviral, antibacterial, or immunostimulating activity in a mammal, said process comprising the steps of:

35 (1) preparing an extract of *Radix Scutellariae*, which comprises the steps of:

- (a) extracting *Radix Scutellariae* with water at a temperature above about 25°C to provide a first extract;
- 5 (b) concentrating the first extract to provide a concentrate having a relative density of from about 1.20 to about 1.25 at 80°C;
- (c) adjusting the pH of the concentrate to a level of from about 1 to about 2 at 80°C;
- 10 (d) cooling the concentrate to produce a precipitate and recovering the precipitate;
- (e) mixing the precipitate with water to obtain a paste;
- (f) adjusting the pH of the paste to about 7;
- 15 (g) dissolving the paste in ethanol to obtain a first ethanolic solution; and optionally filtering the solution to remove any suspended impurities; and
- (h) concentrating the first ethanolic solution to obtain an extract of *Radix Scutellariae* having a relative density less than about 1.03 at 40°C; and
- 20 (2) preparing a combined extract of *Flos Lonicerae* and *Fructus Forsythiae* which comprises the steps of:
- (a) extracting a mixture of *Flos Lonicerae* and *Fructus Forsythiae* with water at a temperature above about 25°C to provide a second extract;
- 25 (b) concentrating the second extract to provide a second concentrate having a relative density of from about 1.20 to about 1.25 at 80°C;
- (c) cooling the second concentrate to about 40°C;
- 30 (d) adding ethanol to the concentrate to obtain a second ethanolic solution and removing any precipitate from the solution;
- (e) concentrating the second ethanolic solution to remove the ethanol and recovering the combined extract of *Flos Lonicerae* and *Fructus Forsythiae*; and

(3) combining the extract of *Radix Scutellariae* and the combined extract of *Flos Lonicerae* and *Fructus Forsythiae* to obtain the RFF extract.

5 20. A process for isolating baicalin in a purified form from a plant material containing baicalin comprising:

(a) contacting said plant material with water at an elevated temperature to produce a water extract;

(b) recovering said water extract;

10 (c) adjusting the pH of said water extract to an acidic pH;

(d) diluting said extract with water to obtain a precipitate;

15 (e) dissolving said precipitate in an aqueous alkali solution to obtain a neutral solution of said precipitate;

(f) diluting said neutral solution with an alcohol to obtain an alcoholic solution;

(g) adjusting the pH of the alcoholic solution to an acidic pH and obtaining a precipitate containing baicalin;

20 and

(h) recovering the purified baicalin.

21. A process for isolating chlorogenic acid in a purified form from a plant material containing chlorogenic acid comprising:

25 (a) contacting said plant material with water to obtain a water extract;

(b) partitioning said water extract with an organic solvent to obtain an organic phase containing chlorogenic acid;

30 (c) concentrating the organic phase to obtain a concentrate;

(d) purifying the concentrate on a chromatographic column; and

35 (e) recovering the purified chlorogenic acid.

22. A process for isolating forsythiaside in a purified form from a plant material containing forsythiaside comprising:

- 5 (a) contacting said plant material with water at an elevated temperature to obtain a water extract;
- (b) concentrating said water extract to obtain a concentrate;
- (c) contacting said extract with an organic solvent to obtain an organic solution;
- 10 (d) recovering and concentrating said organic solution to obtain a residue containing forsythiaside;
- (e) purifying said residue on a chromatographic column; and
- (f) recovering the purified forsythiaside.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/00334

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K35/78 C07H15/26 C07H3/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE WPI Section Ch, Week 9713 Derwent Publications Ltd., London, GB; Class B04, AN 97-133214 XP002101772 & CN 1 089 848 A (HARBIN NO 4 TRADITIONAL CHINESE PHARM FA), 27 July 1994 see abstract	1-22
X	EP 0 348 509 A (TSUMURA & CO) 3 January 1990 see page 2, line 1 - line 19	1-4, 8-11, 20
X	WO 97 25054 A (LI HONG ;WU LETING (CN)) 17 July 1997 ABSTRACT	1-22
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

3 May 1999

Date of mailing of the international search report

25/05/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040. Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Rempp, G

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/00334

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE WPI Section Ch, Week 9743 Derwent Publications Ltd., London, GB; Class B04, AN 97-458038 XP002101773 & CN 1 117 851 A (SHANDONG COLLEGE TRADITIONAL CHINESE MED), 6 March 1996 see abstract</p>	1-22
X	<p>DATABASE WPI Section Ch, Week 8603 Derwent Publications Ltd., London, GB; Class B05, AN 86-018507 XP002101774 & JP 60 243016 A (TSUMURA JUNTENDO KK) , 3 December 1985 see abstract</p>	1-22

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/ 00334

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 9-14, 17, 18
because they relate to subject matter not required to be searched by this Authority, namely:

see FURTHER INFORMATION sheet PCT/ISA/210
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 99 00334

FURTHER INFORMATION CONTINUED FROM PCT/SA/ 210

Although claims 9-14,17,18 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

Claims Nos.: 9-14,17,18

Rule 39.1(iv) PCT - Method for treatment of the human or animal body by therapy

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/00334

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0348509 A	03-01-1990	JP 1835115 C	11-04-1994
		JP 2069415 A	08-03-1990
		JP 1153646 A	15-06-1989
		JP 1784069 C	31-08-1993
		JP 4070287 B	10-11-1992
		JP 1163120 A	27-06-1989
		JP 1836443 C	11-04-1994
		JP 1163121 A	27-06-1989
		JP 1836444 C	11-04-1994
		JP 1163122 A	27-06-1989
		JP 1836445 C	11-04-1994
		AT 87204 T	15-04-1993
		DE 3879688 A	29-04-1993
		WO 8905141 A	15-06-1989
WO 9725054 A	17-07-1997	CN 1154246 A	16-07-1997
		AU 1364497 A	01-08-1997